

# ICU

## MANAGEMENT & PRACTICE

THE OFFICIAL MANAGEMENT JOURNAL OF ISICEM

VOLUME 16 - ISSUE 4 - WINTER 2016



# Quality

### PLUS

Biomarkers in Heart Failure

Difficult Intubation

Emergency Intraosseous Access

Pain Assessment and Management

Dexmedetomidine, Delirium & Sleep

Early Mobilisation: From Concept to Reality

What Can Psychologists Do in Intensive Care?

Clinical Pharmacist Role

Social Media: Blessing or Curse?

Learning to Lead an ICU

Renal Replacement Therapy for Acute Kidney Injury

Country Focus: UK





### Clarence Chant

Director of Pharmacy  
Pharmacy Department  
St. Michael's Hospital  
Toronto, Canada

Adjunct Professor  
Leslie Dan Faculty of Pharmacy  
University of Toronto  
University of Toronto

CHANTC@smh.ca



### Norman F. Dewhurst

Clinical Pharmacy Specialist/Leader  
Pharmacy Department  
St. Michael's Hospital  
Toronto, Canada

DewhurstN@smh.ca

# CLINICAL PHARMACIST ROLE IN THE ICU

We provide an overview of the various facets of pharmacist practice in the intensive care unit (ICU), the current extent to which pharmacists are present in the ICU, along with a discussion on barriers and lessons learned in garnering support for such a role.

## Reduce Medication Errors

A landmark study in 1999 reported that pharmacist attendance in ICU rounds reduced the rate of preventable adverse drug events by 66% (Leape et al. 1999). Using the EU study figures, this would extrapolate to over 1200 lives saved every year. Other publications further support improved clinical outcomes due to pharmacist interventions. In a retrospective review of patients with thromboembolic disease, critical care pharmacists were able to significantly reduce patient mortality, ICU length of stay, bleeding complications, and need for blood product transfusions (MacLaren and Bond 2009). Recently, the PROTECTED-UK study involving 21 ICUs over 2 weeks reported that pharmacists reviewed 20,517 medication orders, 3,294 (16.1%) of which required interventions to optimise medication therapy (Shulman et al. 2015). Of the interventions, the majority (87.7%) were accepted by the prescriber, 6.8% were medication errors and 66% were deemed to be high risk in nature. Other studies have reported estimated cost-savings or avoidance of \$1.7-2.1 million over a 2 year period, making a return on investment of 7 to 1 (Weant et al. 2009).

## Education, Research, Administration

Critical care pharmacists are a valuable resource in providing education to clinical team members in addition to pharmacist trainees. In a neonatal ICU, a pharmacist-led staff education and risk management programme reduced medication errors from 24.1 to 5.1 per 1000 neonatal activity days (Simpson et al. 2004). Similarly, physician orientation and education were shown to reduce prescribing error rates. A panel consisting of a pharmacist and paediatrician, using a standardised predefined criteria, rated the severity of the errors and found a reduction

in severe errors from 29.7% to 7% (Alagha et al. 2011).

Critical care pharmacists can also lead and/or participate in clinical research. In a Canadian survey specifically on this topic involving 215 pharmacists, 41.4% reported being moderately to highly involved in research (Perreault et al. 2012). Finally, pharmacists can also be involved in more administrative/leadership type roles, such as quality improvement. In one pharmacist-driven quality improvement initiative (QI), an interdisciplinary protocol was shown to significantly improve process measure compliance with spontaneous awakening trials from a baseline of 20% to 97-100%, which was sustained 8 months following the programme (Stollings et al. 2015).

It would appear that there is an abundance of literature demonstrating ICU pharmacist ability to improve financial, clinical, and process outcomes. It is therefore disheartening to observe that 17 years after the publication of the landmark study (Leape et al. 1999), adoption is far less than 100%, despite wide support by professional organisations and patient safety experts (MacLaren et al. 2006; Brilli et al. 2001). A few barriers and lessons learned are presented below as a starting point to assist those contemplating such an undertaking.

## Building the Business Case

For most institutions, in order to obtain a new ICU pharmacist, a convincing business case is required. While specific requirements differ depending on local contexts, this usually involves a needs assessment, an environmental scan of comparator institutions, proposed service model, cost of service (e.g. pharmacist yearly salary and benefits), potential cost savings, and risk-benefit assessment of implementation. An environmental scan can be done locally within

Caring for critically ill patients in an intensive care unit (ICU) is considered a standard of care in today's environment. However, the ICU is a rapidly changing, complex, and costly environment where polypharmacy is the norm and medications are frequently used in combinations involving ever-changing doses based on physiologic responses and critical illness-related organ dysfunction. This creates the 'perfect storm' scenario that is ripe for medication errors. A study over a three-week period in two ICUs in the U.S. found an adverse event rate of 80.5/1000 patient-days, with medications being responsible for 78% of the serious events (Rothschild et al. 2005). This error rate is not an isolated phenomenon; a European study conducted across 27 countries and 113 ICUs involving 1,328 patients revealed that during a brief 24-hour observation period 81% of ICUs reported at least one parenteral medication error that involved 37% of patients (Valentin et al. 2009). This translated into an error rate of 74.5 errors per 100 patient days, with 7 patients experiencing permanent harm and 5 patient deaths due to medication errors. From a cost perspective, medications are the fourth largest contributor to total ICU costs, and account for approximately 38% of total drug costs in a hospital (Weber et al. 2003). Fortunately, the role of pharmacists in reducing medication errors and costs is well established.

the city or health region, published literature, or where available, national data such as the Canadian Hospital Pharmacy report (MacLaren et al. 2006; Hospital Pharmacy in Canada Editorial Board 2015). The caveat is that a significant portion of the overall cost savings made by ICU pharmacists is not in direct drug costs, but in prevention of costs due to errors. This is, albeit very unfortunately, viewed differently by administrators and finance personnel as not 'real dollars saved'. Therefore the 'sales pitch' often needs to centre on quality of care and/or meeting of regulatory or accreditation requirements, supported by any local/national quality agenda/initiative, and preferably in alignment with institution-specific objectives. Failing that, another method to demonstrate the worth of an ICU pharmacist has sometimes come from a trial period where another pharmacist with the appropriate knowledge/skills is redeployed to practise in the ICU for a short period while documenting the interventions made. This type of trial period allows for gathering of local data, which may be more convincing, but perhaps more importantly, allows the ICU care team to witness first-hand the benefits of having a pharmacist. Often the clinical team members (e.g. nurses and physicians) will become the best champions and advocates. Relationship building with the ICU team is a key factor in success, and this may be established through other channels such as collaborative work in a project for the ICU (e.g. computer order entry implementation) or through pharmacotherapy guideline development during Pharmacy and Therapeutics committee participation.

### Education and Training

Training for ICU pharmacotherapy is usually not the focus of many undergraduate pharmacy curricula and ICU clinical rotations/clerkships are often viewed by students as 'difficult to pass' rotations. Therefore, students' interest in ICU as a practice area is not widespread, limiting the qualified recruitment pool when a position is secured. While there are specialty residency programmes in critical care, their availability does not match needs, as demonstrated by the recent survey reporting that only 5.9% of critical care pharmacists have completed a critical care specialty residency (MacLaren et al. 2006). Therefore, finding qualified pharmacists to fill ICU positions is challenging and may result in filling them with less trained personnel, often producing less than optimal acceptance. Fortunately, with dedicated courses in ICU pharmacotherapy appearing in the elective portion of some pharmacy curricula, the addition of board certification in Critical Care Pharmacy by the Board of Pharmacy Specialties in the U.S., more and more training programmes and opportunities will be forthcoming to help close the qualified personnel shortage and needs gap.

### ICU Pharmacist Activities

The Society of Critical Care Medicine and the American College of Clinical Pharmacy published a position paper in 2000 on various activities that can/should be performed by an ICU pharmacist, dividing these activities into fundamental, desirable, and optimal levels (Rudis and Brandl 2000). The list is quite all encompassing, and almost daunting for institutions that

currently don't have such a position. Focusing on part of the fundamental activities, along with meticulous documentation of the interventions/outcomes, using either a homegrown or commercially available tool, should be the initial phase before progressing to desirable or optimal activities. This approach is corroborated by the recent U.S. survey where fundamental activities (e.g. providing drug information) are provided by 83.9% of respondents, desirable activities (e.g. therapeutic management advice to physicians) are performed by 63.8% of respondents, and optimal activities (e.g. ICU research) are performed by 19.5% of respondents (MacLaren et al. 2006).

### Conclusion

In conclusion, while it is unsatisfactory to see that ICU pharmacists are not present in all institutions that have an ICU, even in countries such as the U.S. and Canada where this practice is much more developed, ongoing support from professional organisations, such as the Faculty of Intensive Care Medicine and the Intensive Care Society in the UK, will hopefully continue to challenge the status quo. Indeed, even in developing countries such as Jordan, India and Brazil, studies on the impact of ICU pharmacists are being published (Leblanc et al. 2008; Hisham et al. 2016; Fideles et al. 2015; Aljbouri et al. 2013). Hopefully in the near future, critical care pharmacists will indeed be 'critical' in all ICUs. ■

### Conflict of Interest

Clarence Chant declares that he has no conflict of interest. Norman Dewhurst declares that he has no conflict of interest.

### References

- Alagha HZ, Badary OA, Ibrahim HM et al. (2011) Reducing prescribing errors in the paediatric intensive care unit: an experience from Egypt. *Acta Paediatrica*, 100(10), e169-174.
- Aljbouri TM, Alkhaldeh MS, Abu-Rumman AK et al. (2013) Impact of clinical pharmacist on cost of drug therapy in the ICU. *Saudi Pharm J*, 21(4): 371-4.
- Brilli RJ, Spevitz A, Branson RD et al. (2001) Critical care delivery in the intensive care unit; Defining clinical roles and the best practice model. *Crit Care Med*, 29(10): 2007-19.
- Faculty of Intensive Care Medicine and Intensive Care Society (2013) Core standards for intensive care units. (2013) [Accessed: 13 October 2016]. Available from [ficm.ac.uk/sites/default/files/Core%20Standards%20for%20ICUs%20Ed.1%20\(2013\).pdf](http://ficm.ac.uk/sites/default/files/Core%20Standards%20for%20ICUs%20Ed.1%20(2013).pdf)
- Fideles GM, de Alcântara-Neto JM, Peixoto Júnior AA et al. (2015) Pharmacist recommendations in an intensive care unit: three-year clinical activities. *Rev Bras Ter Intensiva*, 27(2): 149-154.
- Hisham M, Sivakumar MN, Veerasekar G. (2016) Impact of clinical pharmacist in an Indian Intensive Care Unit. *Indian J Crit Care Med*, 20(2): 78-83.
- Hospital Pharmacy in Canada Editorial Board (2015) Hospital pharmacy in Canada 2013-2014 report. [Accessed: 15 August 2016] Available from [lillyhospitalsurvey.ca/hpc2/content/2015\\_report/FULL-2015.pdf](http://lillyhospitalsurvey.ca/hpc2/content/2015_report/FULL-2015.pdf)
- Leape LL, Cullen DJ, Clapp MD, et al. (1999) Pharmacist participation on physician rounds and adverse drug events in the intensive care unit. *JAMA*, 282(3): 267-270.
- LeBlanc JM, Seoane-Vazquez EC, Arbo TC et al. (2008) International critical care hospital pharmacist activities. *Intensive Care Med*, 34(3): 538-42.
- MacLaren R, Bond CA (2009) Effects of pharmacist participation in intensive care units on clinical and economic outcomes of critically ill patients with thromboembolic or infarction-related events. *Pharmacotherapy*, 29(7): 761-8.
- MacLaren R, Devlin JW, Martin SJ et al. (2006) Critical care pharmacy services in United States hospitals. *Ann Pharmacother*, 40: 612-8.
- Perreault MM, Thiboutot Z, Burry L, et al. (2012) Canadian survey of critical care pharmacists' views and involvement in clinical research. *Ann Pharmacother*, 46(9): 1167-73.
- Rothschild JM, Landrigan CP, Cronin JW et al. (2005) The critical care safety study: the incidence and nature of adverse events and serious medical errors in intensive care. *Crit Care Med*, 33(8): 1694-1700.
- Rudis MI, Brandl KM, for the SCCM and ACCP Clinical Pharmacy task force on critical care pharmacy services (2000) Position paper on critical care pharmacy services. *Crit Care Med*, 28(11): 3746-50.
- Shulman R, McKenzie CA, Landa J et al. (2015) Pharmacist's review and outcomes: Treatment-enhancing contributions tallied, evaluated, and documented (PROTECT-UK). *J Crit Care*, 30(4): 808-13.
- Simpson JH, Lynch R, Grant J et al. (2004) Reducing medication errors in the neonatal intensive care unit. *Arch Dis Child Fetal Neonatal Ed*, 89(6): F480-2.
- Stollings JL, Foss JJ, Ely EW, et al. (2015) Pharmacist leadership in ICU quality improvement: Coordinating spontaneous awakening and breathing trials. *Ann Pharmacother*, 49(8): 883-91.
- Valentin A, Capuzzo M, Guidet B et al. (2009) Errors in administration of parenteral drugs in intensive care units: multinational prospective study. *BMJ*, 338: b814.
- Weant KA, Armitstead JA, Ladha AM et al. (2009) Cost effectiveness of a clinical pharmacist on a neurosurgical team. *Neurosurgery*, 65(5): 946-51.
- Weber RJ, Kane SL, Oriolo VA et al. (2003) Impact of intensive care unit (ICU) drug use on hospital costs: a descriptive analysis, with recommendations for optimizing ICU pharmacotherapy. *Crit Care Med*, 31(1 Suppl): S17-S24.